

HERBICIDE EVALUATION IN ARKANSAS COTTON, 2000

Marilyn McClelland

Research Associate

Crop, Soil, and Environmental Sciences Department

Jim Barrentine

Professor and Head

Crop, Soil, and Environmental Sciences Department

Ken Smith

Extension Weed Scientist

Southeast Research and Extension Center

Nilda Burgos

Assistant Professor

Crop, Soil, and Environmental Sciences Department

ACKNOWLEDGMENTS

The following companies provided financial support and chemicals used in the studies: AgrEvo, American Cyanamid, BASF, Cheminova, Dow AgroSciences, DuPont, FMC, Griffin, Helena (Setre), Monsanto, Novartis, Rhone-Poulenc, Riverside Terra, Rohm & Haas, Uniroyal, Valent, and Zeneca.

The assistance of the following individuals is gratefully acknowledged: Larry Earnest, Superintendent, Southeast Branch Station, Rohwer; Vaughn Skinner, Farm Manager, Main Experiment Station, Fayetteville; Michelle Mobley, Graduate Assistant, Main Experiment Station, Fayetteville; Scott Payne, Graduate Assistant, Main Experiment Station, Fayetteville; Oscar Sparks, Graduate Assistant, Main Experiment Station, Fayetteville; Erin Cable, Graduate Assistant, Main Experiment Station, Fayetteville; Ford Baldwin, Extension Weed Specialist, Cooperative Extension Service, Little Rock; Jeff Branson, Research Specialist, Southeast Branch Station, Rohwer; Robin Namenek, Research Specialist, Southeast Branch Station, Rohwer; Claude Kennedy, Director, Cotton Branch Experiment Station, Marianna; James Hornbeck, Research Specialist, Cotton Branch Experiment Station, Marianna; and Marci Milus and Lynn McCoy, secretarial staff.

CONTENTS

Introduction	1
Methods	1
Abbreviations of Terms	2
Tables	
Table 1. Benefits of pyriithiobac (Staple) in Roundup Ready (glyphosate-tolerant) cotton, Fayetteville, 2000.	3
Table 2. Benefits of pyriithiobac (Staple) in Roundup Ready (glyphosate-tolerant) cotton, Marianna, 2000.	8
Table 3. Evaluation of pyriithiobac (Staple) and glyphosate (Roundup Ultra) mixtures, Rohwer, 2000.	11
Table 4. Pyriithiobac (Staple) and glyphosate combinations applied PRE and POST, Rohwer, 2000.	13
Table 5. Glyphosate (Roundup Ultra) Programs, Fayetteville, 2000.	16
Table 6. Glyphosate cotton selectivity trial, Marianna, 2000.	21
Table 7. Value of soil residual herbicides in Roundup Ready (glyphosate-tolerant) cotton, Rohwer, 2000.	25
Table 8. Morningglory control in Roundup Ready (glyphosate-tolerant) cotton, Rohwer, 2000.	28
Table 9. Glyphosate cotton selectivity with different adjuvant systems, Marianna, 2000.	31
Table 10. Cloransulam (FirstRate) and cloransulam + flumetsulam (Frontrow) post-directed on Roundup Ready cotton, Marianna, 2000.	33
Table 11. Prometryn (Caparol) preemergence in glyphosate-tolerant cotton, Rohwer, 2000.	37
Table 12. Fomesafen (Reflex) tank mix in cotton, Marianna, 2000.	39
Table 13. Fomesafen (Reflex) preemergence tank mixtures, Rohwer, 2000.	42
Table 14. CGA-362622 in cotton, Fayetteville, 2000.	44
Table 15. CGA-362622 rates and timings, Rohwer, 2000,	48
Table 16. CGA-362622 sequential applications, Rohwer, 2000.	51

Table 17. CGA-362622 in comparison to pyriithiobac (Staple), Rohwer, 2000.	53
Table 18. CGA 362622 in glyphosate-tolerant cotton, Rohwer, 2000.	56
Table 19. CGA 36622 and glyphosate (Roundup) timings, Rohwer, 2000.	59
Table 20. Touchdown (glyphosate): POST control in glyphosate-tolerant cotton with new formulation (3SL), Fayetteville, 2000.	61
Table 21. Touchdown (3 SL formulation) selectivity on cotton, Marianna, 2000.	65
Table 22. Evaluation of glyphosate-tolerant cotton to Touchdown (3 SL formulation), Rohwer, 2000.	67
Table 23. Weed control in glyphosate-tolerant cotton with Touchdown, Rohwer, 2000.	68
Table 24. Buctril study, Marianna, 2000.	70
Table 25. Cotton programs in BXN (bromoxynil-resistant) cotton, Rohwer, 2000.	73
Table 26. Pigweed control in BXN (bromoxynil-resistant) cotton, Rohwer, 2000.	76
Table 27. Pyriithiobac (Staple) / fluometuron (Cotoran) rates in BXN (bromoxynil-resistant) cotton, Rohwer, 2000.	79
Table 28. CGA 362622 in BXN (bromoxynil-resistant) cotton, Rohwer, 2000.	81
Table 29. Clethodim (Select) in BXN (bromoxynil-resistant) cotton, Rohwer, 2000.	84
Table 30. Post-directed herbicide programs in cotton, Fayetteville, 2000.	86
Table 31. Layby study on cotton, Marianna, 2000.	90
Table 32. Flumioxazin (Valor) for layby and residual weed control, Rohwer, 2000.	94
Table 33. Efficacy of carfentrazone (Aim) post-directed in cotton, Rohwer, 2000.	96
Table 34. Harvade (dimethipin) in combination with residual post-directed herbicides, Rohwer, 2000.	98
Table 35. Harvade (dimethipin) tank mixtures for layby control, Rohwer, 2000.	100
Table 36. Transgenic cotton in ultra-narrow row, Marianna, 2000.	102

Appendix Tables

Appendix Table 1. Common and trade names, formulation (pounds of active ingredient or acid equivalent per gallon), sponsoring companies, and chemical names of herbicides	111
Appendix Table 2. Trade names and common names.	112
Appendix Table 3. Common, Bayer codes, and scientific names of plant species	113
Appendix Table 4. Climatological data, Main Experiment Station, Fayetteville, 2000.	114
Appendix Table 5. Climatological data, Cotton Branch Experiment Station, Marianna, 2000.	115
Appendix Table 6. Climatological data, Southeast Branch Experiment Station, Rohwer, 2000.	116